

Table 2. List of numerical experiments.

Experiment	Description—variation from reference model
EX1	–reference model
EX2	–wet olivine mantle ¹ : $A = 4.89 \times 10^{-15} \text{ Pa}^{-n}/\text{s}$, $n = 3.5$, $Q = 515 \text{ kJ/mol}$ in regions 2, 3, 4, 5.
EX3	–dry olivine mantle ¹ : $A = 4.85 \times 10^{-17} \text{ Pa}^{-n}/\text{s}$, $n = 3.5$, $Q = 535 \text{ kJ/mol}$ in regions 2, 3, 4, 5.
EX4	–increase range of strain for plastic softening: $\epsilon_1=0.5 \rightarrow \epsilon_2 = 5.0$ (for same $\phi_1 = 15^\circ$ and $\phi_2 = 2^\circ$).
EX5	–suppress strain softening: set $\phi_1 = 15^\circ$ and $\phi_2 = 15^\circ$.
EX6	–increase initial geotherm in model (see text).
EX7	–increase crustal thickness to $\sim 40.8 \text{ km}$.
EX8	–make mantle neutrally buoyant by setting $\alpha = 0$ in all regions.
EX9	–decrease imposed convergence velocity to $v_x = 0.5 \text{ cm/yr}$.
EX10	–increase imposed convergence velocity to $v_x = 4.5 \text{ cm/yr}$.

Reference for viscosity parameters: 1. *Hirth and Kohlstedt [1996]*.

:

: